



Use this wallet card to help you remember some of the potential side effects associated with your Pr KEYTRUDA® (pembrolizumab) treatment.

If you experience any symptoms, talk to your medical team without delay.

Carry this wallet card with you at all times

in case you visit an emergency room or have to meet with a healthcare professional who is not part of your KEYTRUDA® treatment team. It will provide them with important information to help understand your treatment and how to deal with your side effects.

My name: _____



For healthcare team in case of emergency:

I am being treated with ^{Pr}KEYTRUDA[®] (pembrolizumab).
KEYTRUDA[®] is not chemotherapy. Adverse events caused
by KEYTRUDA[®] need to be treated urgently and appropriately.

Please contact my oncology team immediately.

My oncologist's name: _____

Office phone: _____

After hours phone: _____

Treated for: _____

What is KEYTRUDA® used for?

KEYTRUDA®, also known as pembrolizumab, is a prescription medicine used to treat:

- A kind of skin cancer called melanoma in adults
 - KEYTRUDA® may be used alone as your first treatment when your skin cancer:
 - has spread or cannot be removed by surgery (advanced melanoma)
- A kind of lung cancer called non-small cell lung cancer in adults
 - KEYTRUDA® may be used alone as your first treatment when your lung cancer:
 - has spread (advanced lung cancer), or
 - has not spread outside your chest (Stage III) and you cannot have surgery or chemotherapy with radiation, and
 - tests positive for “PD-L1”, and
 - if your tumour does not have an abnormal “EGFR” or “ALK” gene

It is not known if KEYTRUDA® is safe and effective in children less than 18 years of age.

People get KEYTRUDA® when their cancer has spread or cannot be taken out by surgery.

Serious side effects and what to do about them

Be aware of possible side effects

Keep this leaflet on you. The symptoms map on the back lists the major side effects you should look out for. These side effects can sometimes become life-threatening and can lead to death. These side effects may happen anytime during treatment or even after your treatment has ended. If you have any of them, call or see your doctor **right away**. Your doctor may give you

other medicines to prevent more severe complications and reduce symptoms. Your doctor may withhold the next dose of KEYTRUDA® or stop treatment with KEYTRUDA®. Do not self-treat any of these symptoms without the approval of your healthcare team.

ALK=anaplastic lymphoma kinase; EGFR=epidermal growth factor receptor; PD-L1=programmed cell death ligand 1

© Merck Sharp & Dohme LLC. Used under license.
© 2015, 2023 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.



Pr **KEYTRUDA**®
(pembrolizumab)

What to look out for



EYES

- Eyesight has changed
- Yellowing of my eyes
- Eye pain
- Dry eyes



MOUTH AND HEAD

- Dry mouth
- Ulcers in my mouth and/or lining of my nose
- Inflammation of the mucous membranes of my mouth
- Cold sores
- Stuffy nose
- Change in my sense of taste
- Faintness or dizziness
- Headaches that will not go away or are unusual
- More thirsty than usual
- Swelling of my face
- Confusion
- Memory problems
- Seizure




THROAT AND CHEST

- New or worse cough
- Ulcers in my throat
- Voice is getting deeper
- Short of breath
- Chest pain
- Rapid heartbeat
- Irregular heartbeat
- Upper respiratory tract infection



SKIN AND HAIR

- Dry skin
- Yellowing of my skin
- Patches of my skin have lost their colour (vitiligo)
- Rash or itchy skin
- Blisters and/or sores on my skin
- Skin is peeling
- Ulcers in my genital area
- Bleeding or bruising more easily than normal
- Sweating more than normal
- Hair is falling out
- Tender lumps on my skin
- Itching



SKIN AND HAIR

- Red skin lesions
- Itchy patches of thick red skin with silvery scales (symptoms of psoriasis)
- Skin conditions resembling acne



STOMACH AND BOWELS

- Less hungry or more hungry than usual
- Nausea or vomiting
- Constipation
- Diarrhea or more bowel movements than usual
- Stools are black, tarry, sticky or have blood or mucus
- Severe stomach pain or tenderness
- Pain on the right side of my stomach
- Swelling of my liver or spleen



URINE

- Changes in the amount of urine
- Need to urinate more often
- Colour of my urine has changed
- Urinary incontinence
- Difficulty urinating



MUSCLES, JOINTS AND LEGS

- Severe or persistent muscle or joint pains
- Muscle problems that can cause weakness and rapid fatigue of muscles or weakness and tingling in arms and legs
- Muscle cramps or spasms
- Swelling in my legs or arms
- Back pain
- Pain, numbness or tingling in my arms or legs or feet or hands



GENERAL

- Fatigue
- Dehydration
- Unusually tired or weak
- Flu-like symptoms
- Fever
- Chills
- Lost or gained weight
- Feel colder than normal
- Swollen lymph nodes
- Low red blood cell count
- General numbness and weakness



RELATED TO THE INFUSION

- Shortness of breath
- Itching or rash
- Dizziness or fever
- Wheezing
- Flushing
- Feeling like passing out

If you have a symptom or side effect that is not listed here, write it down in your diary included in this package and **tell your healthcare team immediately.** Do not wait until your next appointment.